

CORRESPONDENCE

use of air as part of the respiratory gas mixture. Increased inspired carbon dioxide may be due to the presence of carbon monoxide or the presence of carbon dioxide due to carbon dioxide absorbent exhaustion or leaking valves in a circle system. If 500 ppm (0.05%) carbon monoxide was present in a patient's breathing circuit and was displayed as an increase of either 0.05% nitrogen or 0.05% inspired carbon dioxide, I speculate that this increase would not be distinguishable from innocuous fluctuations of these gases. Therefore, I suggest that, before direct detection of carbon monoxide by mass spectrometry can be used to warn of a patient's exposure to carbon monoxide during anesthetic breakdown, studies must be conducted to show the validity of this technique with clinically relevant concentrations of carbon monoxide.

Harvey J. Woehlck, M.D.
Department of Anesthesiology

Anesthesiology
83:1384-1385, 1995
© 1995 American Society of Anesthesiologists, Inc.
Lippincott-Raven Publishers

Binding of Halothane to Serum Albumin: Relevance to Theories of Narcosis

To the Editor:—The report by Johansson *et al.* provides further insight into the molecular site at which general anesthetics act.¹ The investigators found that halothane quenches the tryptophan fluorescence of bovine serum albumin in a concentration-dependent manner with a dissociation constant of 1.8 mM. They also reported that diethyl ether competes with halothane with a 50% inhibition concentration of 39 mM.

These concentrations surpass those required for anesthesia. At 1.8 mM, halothane equals a partial pressure at 37°C of 0.06 atm ($1.8 \text{ mM} = (1.8 \times 10^{-6} \text{ mol/ml}) (2.5436 \times 10^4 \text{ ml/mol})/0.75$, where 0.75 is the partition coefficient for halothane in Krebs' solution at 37°C.² This exceeds the anesthetizing partial pressure of halothane in humans by a factor of 8.³ Similarly, the partial pressure of ether at 39 mM equals 0.76 atm, assuming a partition coefficient of 13.⁴ This exceeds the anesthetizing partial pressure of ether in humans by a factor of 40.⁵

Although Johansson *et al.* performed their studies at 25°C, the above ratios (8 for halothane and 40 for ether) for 37°C will approximate ratios at 25°C because of the counterbalancing changes in solubility and potency of anesthetics with decreasing temperature.⁶ Furthermore, the dissociation constant of 1.8 mM for halothane quenching found by Johansson *et al.* is also an order of magnitude greater than the anesthetic potency of halothane measured in animals at lower temperatures: The righting reflex EC_{50} of halothane in tadpoles is approximately 0.1 mM at 20°C.⁷ The calculations of partial pressure also assume that the solution used in the experiment was equivalent to an isotonic salt solution. If the albumin added appreciably to the solubility of halothane, this would lower the partial pressure calculated for halothane but not that for ether, whose solubility in blood scarcely differs from that in water.^{8,9}

Froedtert Memorial Lutheran Hospital
9200 West Wisconsin Avenue
Milwaukee, Wisconsin 53226

References

1. Stewart RD: The effect of carbon monoxide on humans. *J Occupational Med* 18:304-309, 1976
2. Breen PH, Isserles SA, Westley J, Roizen MF, Taitelman UZ: Combined carbon monoxide and cyanide poisoning: A place for treatment? *Anesth Analg* 80:671-677, 1995

(Accepted for publication September 18, 1995.)

Even allowing for these factors, it appears that the partial pressures applied exceed those that produce anesthesia. If so, can the results provide us with insights into mechanisms of anesthetic action? Does the five-fold difference in the ratios for ether and halothane (8 *vs.* 40) mean that the finding for halothane does not apply equally to all anesthetics, and thus that the tryptophan site is not representative of a relevant anesthetic site of action? Finally, do results obtained at 25°C apply at the higher temperatures sustained by homeotherms?

Edmond I. Eger II, M.D.
Professor and Vice Chairman for Research
Donald D. Koblin, Ph.D., M.D.
Professor
Department of Anesthesia
University of California, San Francisco
Box 0464, Science 455
513 Parnassus Avenue
San Francisco, California 94143-0464

References

1. Johansson JS, Eckenhoff RG, Dutton PL: Binding of halothane to serum albumin demonstrated using tryptophan fluorescence. *ANESTHESIOLOGY* 83:316-324, 1995
2. Renzi F, Waud BE: Partition coefficients of volatile anesthetics in Krebs' solution. *ANESTHESIOLOGY* 47:62-63, 1977
3. Saidman LJ, Eger EI II: Effect of nitrous oxide and of narcotic premedication on the alveolar concentration of halothane required for anesthesia. *ANESTHESIOLOGY* 25:302-306, 1964

CORRESPONDENCE

4. Eger EI II, Shargel RO, Merkel G: Solubility of diethyl ether in water, blood and oil. *ANESTHESIOLOGY* 24:676-678, 1963

5. Saidman LJ, Eger EI II, Munson ES, Babad AA, Muallem M: Minimum alveolar concentrations of methoxyflurane, halothane, ether and cyclopropane in man: Correlation with theories of anesthesia. *ANESTHESIOLOGY* 28:994-1002, 1967

6. Eger EI II, Saidman LJ, Brandstater B: Temperature dependence of halothane and cyclopropane anesthesia in dogs: Correlation with some theories of anesthetic action. *ANESTHESIOLOGY* 26:764-770, 1965

7. Firestone S, Firestone L, Ferguson C, Blanck D: Staurosporine, a protein kinase C inhibitor, decreases the general anesthetic requirement in *Rana pipiens* tadpoles. *Anesth Analg* 77:1026-1030, 1993

8. Eger EI II, Larson CP Jr: Anaesthetic solubility in blood and tissues: Values and significance. *Br J Anaesth* 36:140-149, 1964

9. Larson CP Jr, Eger EI II, Severinghaus JW: The solubility of halothane in blood and tissue homogenates. *ANESTHESIOLOGY* 23:349-355, 1962

(Accepted for publication September 18, 1995.)

Anesthesiology
83:1385, 1995

© 1995 American Society of Anesthesiologists, Inc.
Lippincott-Raven Publishers

In Reply:—Eger and Koblin raise several issues concerning the validity of albumin as a model to study mechanisms of inhalational anesthetic action and the relevance of the concentrations both of halothane and diethyl ether used in our studies. The principal aim of our communication¹ was to describe an alternative and useful approach to determining anesthetic-protein interactions. Albumin was selected to compare the tryptophan fluorescence quenching method with the previous studies on anesthetic binding to this protein, which used ¹⁹F-NMR² and photoaffinity labeling.³ Our experiments were performed at 25°C to permit comparison with the earlier studies^{2,3} conducted at similar temperatures and because the direct information available⁴ suggests that, at least over the temperature range from 5°C to 20°C, volatile general anesthetic binding to protein targets only changes by 20–30%.

The measured dissociation constants of halothane and diethyl ether determined using quenching of tryptophan fluorescence¹ are greater than those required to maintain the anesthetic state in animals. It would be remarkable if the anesthetic binding sites on albumin exactly reproduced the clinical pharmacology of the inhaled general anesthetics. However, the fact that these tryptophan domains bind anesthetics within an order of magnitude of their clinical ED₅₀'s suggests that they may share some characteristics with the physiologically relevant sites. Although this was not the primary goal of the work, this is a satisfactory first approximation, because a tenfold change in affinity only corresponds to a binding energy of 1.4 kcal/mol (comparable to a single hydrogen bond⁵). Our study opens the door methodologically for a molecular description of anesthetic binding sites in proteins: For example, one avenue would be the use of site-directed mutagenesis of the albumin sites to conduct a structure-affinity study with a variety of volatile anesthetic molecules.

In further response to the points made by Eger and Koblin concerning the weakness of the binding and the failure to reproduce the relative order of potencies seen clinically, it should be emphasized that it remains to be established that these structurally diverse molecules act at a single locus. Furthermore, it is far from clear that clinical EC₅₀ values must correspond directly to experimental dissociation constants. These questions remain unanswered, because

the methodology to directly study volatile anesthetic binding has been unavailable until recently and is under development. Binding studies accounting for anesthetic desolvation and the formation of interactions with protein sites with a focus on molecular structure are needed. Using our approach¹ and other techniques,^{2,3} the answers to these questions are forthcoming.

Jonas S. Johansson, M.D., Ph.D.
Roderic G. Eckenhoff, M.D.
P. Leslie Dutton, F.R.S.

Departments of Anesthesiology and
the Johnson Research Foundation
Department of Biochemistry and Biophysics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

References

1. Johansson JS, Eckenhoff RG, Dutton PL: Binding of halothane to serum albumin demonstrated using tryptophan fluorescence. *ANESTHESIOLOGY* 83:316-324, 1995
2. Dubois BW, Cherian SF, Evers AS: Volatile anesthetics compete for common binding sites on bovine serum albumin: A ¹⁹F-NMR study. *Proc Natl Acad Sci USA* 90:6478-6482, 1993
3. Eckenhoff RG, Shuman H: Halothane binding to soluble proteins determined by photoaffinity labeling. *ANESTHESIOLOGY* 79:96-106, 1993
4. Dickinson R, Franks NP, Lieb WR: Thermodynamics of anesthetic/protein interactions: Temperature studies on firefly luciferase. *Biophys J* 64:1264-1271, 1993
5. Shirley BA, Stanssens P, Hahn U, Pace CN: Contribution of hydrogen bonding to the conformational stability of ribonuclease T1. *Biochemistry* 31:725-732, 1992

(Accepted for publication September 18, 1995.)